

# 2019 VCE Mathematical Methods 1 (NHT) examination report

## Specific information

This report provides sample answers or an indication of what answers may have been included. Unless otherwise stated, these are not intended to be exemplary or complete responses.

### Question 1a.

$$y = 2e^x - e^{-x} \text{ so } \frac{dy}{dx} = 2e^x + e^{-x}$$

$$\text{Or } \frac{dy}{dx} = \frac{4e^{2x}e^x - (2e^{2x} - 1)e^x}{(e^x)^2} = \frac{2e^{3x} + e^x}{e^{2x}} \text{ (quotient rule)}$$

Some students used a combination of product and chain rules.

### Question 1b.

$$f'(x) = 2x \cos(3x) - 3x^2 \sin(3x)$$

$$f'\left(\frac{\pi}{3}\right) = -\frac{2\pi}{3}$$

### Question 2

$$f(x) = \frac{2}{3}x^3 - \frac{3}{4}x^{\frac{1}{3}} + c$$

$$\text{where } c = f(1) - \frac{2}{3} + \frac{3}{4} = -\frac{7}{4} - \frac{2}{3} + \frac{3}{4} = -\frac{5}{3}$$

$$\text{So } f(x) = \frac{2}{3}x^3 - \frac{3}{4}x^{\frac{1}{3}} - \frac{5}{3}$$

### Question 3a.

$$\int_2^7 \frac{1}{x - \sqrt{3}} dx = \left[ \log_e(x - \sqrt{3}) \right]_2^7 = \log_e \left( \frac{7 - \sqrt{3}}{2 - \sqrt{3}} \right)$$

$$\int_2^7 \frac{1}{x + \sqrt{3}} dx = \left[ \log_e(x + \sqrt{3}) \right]_2^7 = \log_e \left( \frac{7 + \sqrt{3}}{2 + \sqrt{3}} \right)$$

**Question 3b.**

$$\begin{aligned} & \frac{1}{2} \left( \frac{1}{x-\sqrt{3}} + \frac{1}{x+\sqrt{3}} \right) \\ &= \frac{1}{2} \left( \frac{(x+\sqrt{3}) + (x-\sqrt{3})}{x^2-3} \right) \\ &= \frac{x}{x^2-3} \end{aligned}$$

**Question 3c.**

$$\begin{aligned} & \int_2^7 \frac{x}{x^2-3} dx \\ &= \frac{1}{2} \int_2^7 \frac{1}{x-\sqrt{3}} + \frac{1}{x+\sqrt{3}} dx \\ &= \frac{1}{2} \log_e \left( \frac{(7-\sqrt{3})(7+\sqrt{3})}{(2-\sqrt{3})(2+\sqrt{3})} \right) \\ &= \frac{1}{2} \log_e(46) \end{aligned}$$

**Question 4a.**

$$g(x) = \log_e(x-3) + 2$$


Domain:  $x > 3$  or  $(3, \infty)$

Range:  $R$

**Question 4bi.**

$$g'(x) = \frac{1}{x-3}$$

Using  $g(4) = 2$  and  $g'(4) = 1$  the tangent is  $y = x - 2$

**Question 4bii.****Question 5a.**

$$\left[ (h(x))^2 \right] = 1 \text{ so } h(x) = 1 \text{ or } -1$$

$$\sqrt{2x+3} - 2 = 1, -1$$

$$\sqrt{2x+3} = 3, 1$$

$$2x+3 = 9, 1$$

$$x = -1, 3$$

Both values are in the domain of  $h$ .

**Question 5b.**

$$\text{Let } y = h^{-1}(x)$$

$$x = \sqrt{2y+3} - 2$$

$$x+2 = \sqrt{2y+3}$$

$$(x+2)^2 = 2y+3$$

$$y = \frac{1}{2}(x+2)^2 - \frac{3}{2}$$

$$\text{Hence } h^{-1}(x) = \frac{1}{2}(x+2)^2 - \frac{3}{2}$$

Domain:  $[-2, \infty)$

**Question 6a.**

Since first two tosses are heads, required probability is

$\text{Pr}(2 \text{ heads out of next } 3) + \text{Pr}(3 \text{ heads out of next } 3)$

$$= \binom{3}{2} \left(\frac{1}{2}\right)^3 + \binom{3}{3} \left(\frac{1}{2}\right)^3 = \frac{3+1}{8} = \frac{1}{2}$$

Use of a tree diagram or any other appropriate method was accepted.

**Question 6b.**

$$\begin{aligned} & \left( \frac{2}{3} - \frac{33}{20} \sqrt{\frac{2}{3} \times \frac{1}{3} \times \frac{1}{18}}, \frac{2}{3} + \frac{33}{20} \sqrt{\frac{2}{3} \times \frac{1}{3} \times \frac{1}{18}} \right) \\ &= \left( \frac{2}{3} - \frac{33}{20} \sqrt{\frac{1}{9^2}}, \frac{2}{3} + \frac{33}{20} \sqrt{\frac{1}{9^2}} \right) \\ &= \left( \frac{29}{60}, \frac{51}{60} \right) \end{aligned}$$

**Question 7a.**

$$\begin{aligned} A(a) &= \int_0^a (\sin(\pi x) - \sin(\pi a)) dx \\ &= \left[ -\frac{\cos(\pi x)}{\pi} - x \sin(\pi a) \right]_0^a \\ &= \frac{-\cos(\pi a) + 1}{\pi} - a \sin(\pi a) \\ &= \frac{1}{\pi} - \frac{1}{\pi} \cos(a\pi) - a \sin(a\pi) \end{aligned}$$

**Question 7b.**

$$A(1) = \frac{2}{\pi}, \quad A\left(\frac{3}{2}\right) = \frac{1}{\pi} + \frac{3}{2}$$

$$\text{Range: } \left[ \frac{2}{\pi}, \frac{2+3\pi}{2\pi} \right]$$

**Question 7ci.**

$$\begin{aligned} \text{Area} &= \int_0^{\frac{4}{3}} (2 \sin(\pi x) + \sqrt{3}) dx \\ &= 2 \int_0^{\frac{4}{3}} \left( \sin(\pi x) + \frac{\sqrt{3}}{2} \right) dx \\ &= 2 \int_0^{\frac{4}{3}} \left( \sin(\pi x) - \sin\left(\frac{4\pi}{3}\right) \right) dx \\ &= 2A(a) \text{ with } a = \frac{4}{3} \end{aligned}$$

Or observe that required area is a dilation by factor 2 of the original area, width  $a = \frac{4}{3}$

**Question 7cii.**

$$A\left(\frac{4}{3}\right) = \frac{4}{\sqrt{3}} + \frac{3}{\pi}$$

$$= \frac{4\pi\sqrt{3} + 9}{3\pi}$$

**Question 8a.**

$$\Pr(W = k) = \binom{50}{k} \left(\frac{1}{6}\right)^k \left(\frac{5}{6}\right)^{50-k}$$

**Question 8b.**

$$\frac{\Pr(W = k + 1)}{\Pr(W = k)} = \frac{\binom{50}{k+1} \frac{1}{6}^{k+1} \times \frac{5}{6}^{49-k}}{\binom{50}{k} \frac{1}{6}^k \times \frac{5}{6}^{50-k}}$$

$$= \frac{(k) \times (50 - k)!}{(k + 1) \times (49 - k)!} \times \frac{1}{6} \times \frac{6}{5}$$

$$= \frac{(50 - k)}{5(k + 1)}$$

**Question 8c.**

$$\Pr(W = k + 1) < \Pr(W = k)$$

$$(50 - k) > 5(k + 1)$$

$$k > \frac{45}{6} \quad \left(\frac{15}{2}\right)$$

Hence

$$\Pr(W = 7) < \Pr(W = 8)$$

$$\Pr(W = 9) < \Pr(W = 8)$$

So greatest for  $k = 8$

Or by argument from features of binomial distribution.